

Export of Treated Produce

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with eChem Thidiazuron 500 SC Liquid. If you are growing produce for export, please check with eChem (Australia) Pty Ltd for the latest information on MRLs and import tolerances BEFORE using eChem Thidiazuron 500 SC Liquid.

Mixing

Add the required amount of eChem Thidiazuron 500 SC Liquid straight into the partly filled spray tank or mixing vat with agitator running, followed by D-C-TRON Cotton Spray Oil, then add the rest of the water. Keep mixture agitated while spraying. Use only freshly prepared sprays. Do NOT store spray mixture overnight.

When tank mixing with emulsifiable concentrates (ECs), add the eChem Thidiazuron 500 SC Liquid to the half filled spray tank or mixing vat and agitate thoroughly. With agitator running, then add the EC product, followed by the D-C-TRON Cotton Spray Oil. Keep mixture agitated while spraying.

Application

Apply as a uniform spray ensuring thorough coverage of the whole plant.

Equipment Aircraft

Apply by aircraft in 20 to 40 L/ha spray mixture and ensure thorough coverage. A steady cross wind is recommended for best penetration. Do not exaggerate swath width.

Ground Sprayers

Standard boom sprays are recommended and must be fitted with mechanical or by-pass agitation. Use sufficient water to obtain complete coverage of all leaves, ie at least 90 to 100 litres of water per hectare. Good coverage should be obtained with three or more nozzles per row depending upon crop height and canopy density. A nozzle configuration including droppers will give maximum coverage in dryland crops with particular care being taken to ensure thorough but even coverage. Overdosing on some plant parts, eg., nozzle overlap along the sides of plants may cause localised leaf freezing. If band spraying dryland crops, care should be taken when calculating the actual applied rate of eChem Thidiazuron 500 SC Liquid per sprayed hectare.

Compatibility

eChem Thidiazuron 500 SC Liquid is compatible in tank mixtures with ethephon, dimethoate and parathion.

CAUTION: When tank mixed with glyphosate, a higher proportion of cotton leaf may be retained (freezing). Do NOT use this mixture on stressed cotton, particularly drought affected cotton.

PRECAUTIONS

Do not use human flagging unless flaggers are protected by engineering controls such as vehicles with cabs.

Re-entry Period

Do not allow entry into treated areas until spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT allow spray drift to contact trees or crops other than cotton, as this product may injure or defoliate other crops, particularly citrus, grapevines and lettuce.

To help reduce drift damage potential:

DO NOT apply eChem Thidiazuron 500 SC Liquid by air when citrus in flush is within 8 km downwind from the point of application.

DO NOT apply eChem Thidiazuron 500 SC Liquid by air when within 800 m of lettuce.

In addition:

DO NOT plant lettuce or carrots in cotton fields where eChem Thidiazuron 500 SC Liquid has been applied within the previous 9 months.

DO NOT plant legumes, lucerne, other root crops or leafy vegetables (besides those noted above) in cotton fields where eChem Thidiazuron 500 SC Liquid has been applied within the previous two months.

DO NOT plant small grains, sorghum or corn in cotton fields where eChem Thidiazuron 500 SC Liquid has been applied within the previous two weeks.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length PVC gloves. After each day's use, wash gloves and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre . Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet available from eChem (Australia) Pty Ltd.

EXCLUSION OF LIABILITY

This product as supplied is of a high grade and suitable for the purpose for which it is expressly intended and must be used in accordance with the directions. The user must monitor the performance of any product as climatic, geographical or biological variables and/or developed resistance may affect the results obtained. No responsibility is accepted in respect of this product, save for those non-excludable conditions implied by the Trade Practices Act or any State or Federal legislation.

D-C-TRON® is a Registered Trademark of Caltex Australia Petroleum Pty. Ltd.

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

eChem

THIDIAZURON 500 SC

LIQUID COTTON DEFOLIANT

ACTIVE CONSTITUENT: 500g/L THIDIAZURON

For the defoliation of cotton prior to harvest.

APVMA APPROVAL NO: 55963/1208

THIS LEAFLET IS PART OF THE LABEL

eChem (Australia) Pty Ltd
Level 4, Lantos Place 80 Stamford Road Indooroopilly, QLD 4068

Ph: 1300 781 649 Fax: 1300 781 650
admin@echem.com.au www.echem.com.au

DIRECTIONS FOR USE (For use in Qld, NSW, WA only)

Restrains

DO NOT apply eChem Thidiazuron 500 SC Liquid when night temperatures fall below 12° C.

DO NOT apply eChem Thidiazuron 500 SC Liquid following application of sodium chlorate, magnesium chlorate or diquat.

CROP/SITUATION	DEFOLIATION CONDITIONS	RATE/HA	CRITICAL COMMENTS
Cotton Defoliation - first or second spray	IDEAL Conditions are hot to warm (30 to 35°C daytime temperature) and the crop is ready to be defoliated with natural senescence already occurring. Nitrogen and water are almost depleted and there is little or no top growth. A good fruit load is set.	50 to 100 mL plus the recommended label rate of D-C-TRON® Cotton Spray Oil or other spray oil of equivalent specifications registered for cotton	Apply the first spray to cotton, which has been grown under good (non-stressed) conditions and is physiologically mature ie., when the top harvestable boll is mature. Application made prior to boll maturity may result in yield and quality reductions. <ul style="list-style-type: none">Use the lower rates of 50 to 75 mL/ha only under warm early season conditions.Use the higher rate for denser canopies or to provide a faster response.A second spray 7 to 10 days later of either eChem Thidiazuron 500 SC Liquid or an alternative harvest aid may be required.
	GOOD Conditions are warm (25 to 30°C daytime temperature) and the crop is ready to be defoliated. Nitrogen and water are low and there is little or no new growth and a good fruit load is set.	100 to 150mL plus the recommended label rate of D-C-TRON Cotton Spray Oil or other spray oil of equivalent specifications registered for cotton	Apply as above, but use these rates under good conditions.
	AVERAGE Conditions are: (i) warm but night temperatures are dropping to 12°C (not below) and the crop is physiologically ready to defoliate or; (ii) temperature is still hot/warm but the crop is not physiologically ready to senesce.	150 to 200 mL plus the recommended label rate of D-C-TRON Cotton Spray Oil or other spray oil of equivalent specifications registered for cotton	Apply as above, but use these rates under average conditions.
	COLD Night temperatures fall below 12°C and/or mean day/night temperatures fall below 18°C.		DO NOT apply eChem Thidiazuron 500 SC Liquid SC under cold conditions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

Cotton processing - NOT REQUIRED WHEN USED AS DIRECTED.

Grazing/Cotton Trash - DO NOT GRAZE TREATED AREAS OR FEED COTTON TRASH TO LIVESTOCK.