

# POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



eChem

# Thiram 600

FUNGICIDE

ACTIVE CONSTITUENT: 600g/L THIRAM

**GROUP M3 FUNGICIDE**

For use as a seed treatment to control various fungal diseases of Chickpeas, Lupins and Sorghum; and as a drench to control fungal diseases of Turf as specified in the Directions for Use table.

20  
Litres

  
**eChem**  
crop protection

## DIRECTIONS FOR USE

### Restrains:

DO NOT mix this product in a slurry with chickpea or lupin rhizobial inoculant.

Only treat good quality sound seed as treatment of mechanically damaged or poor quality seed may result in reduced germinations and/or reduced seedling vigour and emergence.

SITUATION	DISEASE	RATE	CRITICAL COMMENTS
TURF	Brown patch ( <i>Rhizoctonia solani</i> ) Helminthosporium leaf spot Fusarium patch ( <i>Fusarium</i> spp.)	160 mL in 20 litres of water per 100m <sup>2</sup>	Apply at first sign of disease and repeat at 7-14 day intervals particularly in humid weather.
	Damping off (not NSW)	200 mL in 60 litres of water per 100m <sup>2</sup>	Apply as a drench to the seedbed. Repeat every 5-7 days during disease susceptible periods.

SEED TREATMENT	DISEASE	RATE	CRITICAL COMMENTS
<b>CHICKPEAS</b> Kabuli (Garbanzo) varieties	Damping off ( <i>Pythium</i> spp) and other soil-borne fungi that cause pre-emergence seed rot and seed-borne <i>Botrytis</i>	2 L/tonne seed	Product can be applied as a slurry by mixing the required volume of eChem Thiram 600 with up to 3 litres of water per tonne of seed. Volume of water used should be sufficient to achieve optimum coverage of the grain (up to 3 litres of water per tonne of seed). Apply only to good quality seed. Move seed up and down a grain auger or apply in a suitable mixer. Thorough mixing of the seed after application is required to ensure good coverage. Handle seed carefully after application and avoid excess wetting to prevent damage to the seed coat.
<b>CHICKPEAS</b> Desi varieties	Seed-borne <i>Botrytis</i>		
<b>LUPINS</b>	Seed-borne <i>Anthraco</i> se	1.7-2 L /tonne seed	

SITUATION	PURPOSE	RATE	CRITICAL COMMENTS
<b>SORGHUM SEED</b> - For Export Only	To meet export requirements	Up to 4 L /tonne seed	Treated seed is for export only. Use of product must be in accordance with importing countries requirements. Be aware of any importing countries residue tolerance limits and do not apply product at a rate that will exceed the tolerance limits of the importing country. Treated seed that is not exported must be disposed of in a manner that does not result in planting or consumption by humans or animals.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD:**

**NOT REQUIRED WHEN USED AS DIRECTED.**

eChem (Australia) Pty Ltd  
Level 4, Lantos Place 80 Stamford Road, Indooroopilly, QLD 4068  
Ph: 1300 781 649 Fax: 1300 781 650  
admin@echem.com.au [www.echem.com.au](http://www.echem.com.au)

## GENERAL INSTRUCTIONS:

### Mixing:

Turf should be in good nutritional condition prior to treatment to prevent damage to sensitive turf species (eg. fescues, bluegrass and turf type rye grass), if repeat applications are made too soon. This product can be applied to seed undiluted or may be mixed with water to achieve optimum coverage of the grain. Volumes of water used will depend upon equipment to be used and product rate. Use only in equipment designed for applying liquids or slurries to seed. As flakes may form on the inside of the container, filter before use. Do not allow product to dry in application equipment as the resulting flakes can block filters and nozzles reducing the performance of the equipment. Clean equipment with water after use.

Shake container well before and either shake or agitate well once mixed with water. Agitate regularly during application.

### Inoculation (Seed):

This product will not affect germination of seed if used as directed, however when rhizobial inoculant is also applied to the seed, the viability of the inoculant may be reduced. Chickpea and lupin inoculant can be applied to the seed in a separate operation after seed treatment. Apply the inoculant on the day of sowing after the fungicide treatment has dried on the seed. Sow into a moist seedbed as soon as possible after treatment. If seed is sown into a dry seedbed the fungicide can cause a reduction in nodulation. Alternatively, inoculant can be sprayed into the furrow at seeding rather than combining with the treated seed. This method of applying inoculant is not recommended in WA.

### Sowing:

Check seed flow through the drill after treatment as treated seed can flow more slowly.

### Storage of Treated Seed:

Store treated seed, if not use immediately after treatment, in a dry well-ventilated place. Poor storage of treated seed such as high moisture combined with high temperature can adversely affect subsequent germination. Treated seed should not be held over from one sowing season to be sowed in subsequent seasons. Treated seed should be stored apart from other seed, seed bags or other containers. Bags should be marked to clearly indicate seed has been treated with this product. Empty bags should not be used for any other purpose.

### Compatibility:

Do not mix with lime Sulphur or Bordeaux sprays. Do not mix with fungicides containing copper.

## Fungicide Resistance Warning

**GROUP M3 FUNGICIDE**

For fungicide resistance management eChem Thiram 600 Flowable Fungicide is a Group M3 fungicide.

Some naturally occurring individual fungi resistant to eChem Thiram 600 and other Group M3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by eChem Thiram 600 or other Group M3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, eChem (Australia) Pty Ltd accepts no liability for any losses that may result from the failure of eChem Thiram 600 to control resistant fungi.

### PROTECTION OF LIVESTOCK:

DO NOT feed grass clippings from treated areas or treated seed to poultry, birds or animals. Seed treated with eChem Thiram 600 must not be used for animal or human consumption. It is important that any treated seed not used for sowing is destroyed.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT:

Dangerous to fish. Do not contaminate streams, rivers or waterways with the chemical or used containers including empty bags which have held treated seed. Do not feed or allow birds or wild animals to access treated seed.

### PRECAUTION:

Do not use treated seed for animal or human consumption. Do not allow treated seed to contaminate grain/other seed intended for animal or human consumption. Do not feed treated seed, or otherwise expose, to wild or domestic birds. Do not re use bags (or containers) that have held seed treated with eChem Thiram 600 for any other purpose.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface

in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product

### SAFETY DIRECTIONS:

May irritate the eyes, nose, throat and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container, preparing the product and using the prepared product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length rubber gloves and face shield or goggles. If product on skin immediately wash area with soap and water. If product in eyes wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

### FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Avoid giving alcohol.

### Material Safety Data Sheet

Additional information is listed in the Material Safety Data Sheet which is available from the supplier.


### Conditions of sale:

eChem (Australia) Pty Ltd Accepts responsibility for the consistent quality of the product however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.

APVMA Approval No.: 65286/50633

BN

DOM



9 328283 100130 1